

This Page Is Inserted by IFW Operations  
and is not a part of the Official Record

## **BEST AVAILABLE IMAGES**

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

**IMAGES ARE BEST AVAILABLE COPY.**

**As rescanning documents *will not* correct images,  
please do not report the images to the  
Image Problem Mailbox.**

**PCT**WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau

## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification 6: <b>C12Q 1/68, C07H 21/04</b>	<b>A1</b>	(11) International Publication Number: <b>WO 98/13526</b> (43) International Publication Date: <b>2 April 1998 (02.04.98)</b>
(21) International Application Number: <b>PCT/US97/17338</b> (22) International Filing Date: <b>26 September 1997 (26.09.97)</b>  (30) Priority Data: 00/026.732      26 September 1996 (26.09.96)      US 08/754,580      21 November 1996 (21.11.96)      US  (71) Applicant (for all designated States except US): <b>OLIGOS ETC. INC. [US/US]: 9775 S.W. Commerce Circle #C-6, Wilsonville, OR 97070 (US).</b>  (71)(72) Applicant and Inventor: <b>WOOLF, Tod. M. [US/US]: 21 Birch Road, Natick, MA 01760 (US).</b>  (72) Inventors; and (75) Inventors/Applicants (for US only): <b>ARROW, Amy [US/US]: 15 Equestrian Ridge Road, Newton, CT 06470 (US). DALE, Roderic, M., K. [US/US]: 26761 S.W. 45th Drive, Wilsonville, OR 97070 (US).</b>  (74) Agents: <b>FRIEBEL, Thomas, E. et al.; Pennie &amp; Edmonds LLP. 1155 Avenue of the Americas, New York, NY 10036 (US).</b>		(81) Designated States: <b>AL, AM, AU, AZ, BA, BB, BG, BR, BY, CA, CN, CU, CZ, EE, GE, GH, HU, ID, IL, IS, JP, KG, KP, KR, KZ, LC, LK, LR, LT, LV, MD, MG, MK, MN, MX, NO, NZ, PL, RO, RU, SG, SI, SK, SL, TJ, TM, TR, TT, UA, US, UZ, VN, YU, ARIPO patent (GH, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).</b>  Published With international search report.

(54) Title: **THREE COMPONENT CHIMERIC ANTISENSE OLIGONUCLEOTIDES**

## (57) Abstract

The present application describes a novel family of oligonucleotide compounds having a novel organization of various modified nucleotides and modified chemical linkages. The application further discloses that limiting the presence and extent of specific modified nucleotides/modified linkages in the oligomers results in enhanced activation of endogenous RNase H activity.

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TC	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KR	Korea	NO	Norway	VU	Vanuatu
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

WO 98/13526

PCT/US97/17338

**THREE COMPONENT CHIMERIC ANTISENSE OLIGONUCLEOTIDES**

This application claims priority to U.S. Applications Nos. 60/026,732 filed September 26, 1996, and 08/754,580 filed November 21, 1996.

**1. FIELD OF THE INVENTION**

This invention relates to antisense oligonucleotides that target mRNAs in cells as substrates for the cellular enzyme RNase H and thereby cause specific degradation of the targeted mRNA. The oligonucleotides have four components: an RNase H activating region; a complementarity region; a 5' end; and a 3' end. The invention optimizes each of the components to resist intracellular nucleases, to increase hybridization to target mRNA, to specifically inactivate target mRNA in cells, and to decrease cytotoxicity.

**2. BACKGROUND TO THE INVENTION**

Antisense polynucleotides are useful for specifically inhibiting unwanted gene expression in mammalian cells. They can be used to hybridize to and inhibit the function of an RNA molecule, typically a messenger RNA, by activating RNase H.

The use of antisense oligonucleotides has emerged as a powerful new approach for the treatment of certain diseases. The preponderance of the work to date has focused on the use of antisense oligonucleotides as antiviral agents or as anticancer agents (Wickstrom, E., Ed., Prospects for Antisense Nucleic Acid Therapy of Cancer and AIDS, New York: Wiley-Liss, 1991; Crooke, S.T. and Lebleu, B., Eds., Antisense Research and Applications, Boca Raton: CRC Press, 1993, pp. 154-182; Baserga, R. and Denhardt, D.T., 1992, Antisense Strategies, New York: The New York Academy of Sciences, Vol. 660; Murray, J.A.H., Ed., Antisense RNA and DNA, New York: Wiley-Liss, 1993).

There have been numerous disclosures of the use of antisense oligonucleotides as antiviral agents. For example, Agrawal et al. report phosphoramidate and phosphorothioate

WO 98/13526

PCT/US97/17338

oligonucleotides as antisense inhibitors of HIV. Agrawal et al., Proc. Natl. Acad. Sci. USA 85:7079-7083 (1988). Zamecnik et al. disclose antisense oligonucleotides as inhibitors of Rous sarcoma virus replication in chicken 5 fibroblasts. Zamecnik et al., Proc. Natl. Acad. Sci. USA 83: 4143-4146 (1986).

The principal mechanism by which antisense oligonucleotides affect a targeted RNA molecule is by activation of the cellular enzyme RNase H, which cleaves the 10 RNA strand of DNA/RNA hybrids. Both phosphodiester and phosphorothioate-linked DNA activates endogenous RNase H, thereby cleaving the targeted RNA (Agrawal, S., et al., Proc. Natl. Acad. Sci. USA 87:1101-5 (1990); Woolf, T.M., et al., Nucleic Acids Res. 18:1763-9 (1990)). However, 15 phosphodiester-linked DNA is rapidly degraded by cellular nucleases and, with the exception of the phosphorothioate-linked DNA, nuclease resistant, non-naturally occurring DNA derivatives do not activate RNase H when hybridized to RNA. While phosphorothioate DNA has the advantage of activating 20 RNase H, phosphorothioate-linked DNA has been associated with non-specific cytotoxic effects and a reduced affinity for RNA (Stein, C.A., et al., Aids Res Hum Retroviruses 5:639-46 (1989); Woolf, T.M., et al., Nucleic Acids Res. 18:1763-9 (1990); Kawasaki, A.M., et al., J. Med. Chem. 36:831-41 25 (1993)).

Chimeric antisense oligos that have a short stretch of phosphorothioate DNA (3-9 bases) have been used to obtain RNase H-mediated cleavage of the target RNA (Dagle, J.M., et al., Nucleic Acids Res. 18:4751-7 (1990); Agrawal, S., et 30 al., Proc. Natl. Acad. Sci. USA 87:1401-5 (1990); Monia, B.P. et al., 1993, J. Biol. Chem. 268:14514). A minimum of 3 DNA bases is required for activation of bacterial RNase H (Putdon, P.J., et al., Nucleic Acids Res. 17:9193-9204; 35 Quartin, R.S., et al., Nucleic Acids Res. 17:7235-7262) and a minimum of 5 bases is required for activation of mammalian RNase H (Monia, B.P., et al., J. Biol. Chem. 268:14514-14522 (1993)). In these chimeric oligonucleotides there is a

WO 98/13526

PCT/US97/17338

central region that forms a substrate for RNase H that is flanked by hybridizing "arms," comprised of modified nucleotides that do not form substrates for RNase H.

Alternatively, extracellular tests using a RNase H-containing HeLa cell extract have been reported wherein the RNase H activating region was placed on the 5' or 3' side of the oligomer. Specifically, these tests reported that a 5' or 3' terminal RNase H activating region composed of phosphodiester 2'-deoxynucleotides joined to a methylphosphonate-linked complementarity region was fully active, but that a 5' terminal RNase H-activating region composed of phosphorothioate 2'-deoxynucleotides joined to a methylphosphonate-linked complementarity region was only partially active. See Col 10, U.S. Pat. No. 5,220,007 to T. Pederson et al..

2'-O-Methyl or 2'-fluoro modified nucleotides have been used for the hybridizing arms of chimeric oligos. Inoue, H., et al., 1987, Nucleic Acids Res. 15:6131-48. The 2'-O-Methyl group increases the affinity of the oligomer for the targeted RNA and increases the activity of the oligomer in cell culture. However, 2'-O-Methyl bases with phosphodiester linkages are degraded by exonucleases and so are not suitable for use in cell or therapeutic applications of antisense. Shibahara, S., et al., 1989, Nucleic Acids Res. 17:239-52. Phosphorothioate 2'-O-Methyl nucleotides are resistant to nucleases as shown in the uniformly phosphorothioate modified oligos described by Monia B.P., et al., 1993, J. Biol. Chem. 268:14514-14522 and terminal phosphorothioate substituted, 2'-O-Methylribo-oligonucleotides, Shibahara, S., et al., 1989, Nucleic Acid Res. 17:239-252. However, fully phosphorothioate substituted oligomers may cause non-specific effects including cell toxicity. Stein, C.A., et al., 1989, Aids Res. Hum. Retrov. 5:639-646; Woolf, T.M., et al., 1990, Nucleic Acids Res. 18:1763-69; Wagner, R.W., 1995, Antisense Res. Dev. 5:113-115; Krieg, A.M., & Stein, C.A., 1995, Antisense Res. Dev. 5:241.

WO 98/13526

PCT/US97/17338

The effects of 2'-Fluoro-oligonucleotides on bacterial RNase H are discussed in Crooke, S.T. et al., 1995, Bioch. J. 312:599-608 and Iwai, S. et al., 1995, FEBS Lett (Neth.) 368:315-20.

- 5 Several other chemistries have been used to make the "arms" or regions of a chimeric oligomer that are not substrates for RNase H. The first chimeric oligomers used methylphosphonate or phosphoramidate linkages in the arms (Dagle, J.M., Walder, J.A. & Weeks, K.L., Nucleic Acids Res. 18:1751-7 (1990); Agrawal, S., et al., Proc. Natl. Acad. Sci. USA 87:1401-5 (1990)). While these compounds functioned well in buffered systems and *Xenopus* oocytes, the arms decreased the hybrid affinity. This decrease in affinity dramatically reduced the antisense activity of the chimeric oligomers in
- 15 mammalian cell culture.

- A number of studies have been reported for the synthesis of ethylated and methylated phosphotriester oligonucleotides and their physico-chemical and biochemical evaluation. Dinucleotides with methyl and ethyl triesters were shown to
- 20 possess greater affinity for polynucleotides possessing complementary sequences (Miller, P.S., et al., J. Am. Chem. Soc. 93:6657, (1971)). However, a few years ago, another group reported lack of, or relatively poor, binding affinity of a heptaethyl ester of oligothymidine with complementary
- 25 polynucleotides (Pless, R.C., and Ts'O, P.O.P., Biochemistry 16:1239-1250 (1977)). Phosphate methylated (P-methoxy) oligonucleotides were synthesized and found to possess resistance towards endonuclease digestion (Gallo, K.L., et al. Nucl. Acid Res. 18:7405 (1986)). A P-methoxy 18-mer
- 30 oligonucleotide was shown to have high T<sub>m</sub> value in duplexes with natural DNA and blocked the DNA replication process at room temperature (Moody, H.M., et al., Nucl. Acid Res. 17:4769-4782 (1989)). Moody et al. concluded that phosphate ethylated (P-ethoxy) oligonucleotides would have poor
- 35 antisense properties.

P-methoxy dimers of DNA bases were synthesized using FMOC as transient protecting group for the exocyclic amino

WO 98/13526

PCT/US97/17338

groups (Koole, L.H., et al., J. Org. Chem. 54:1657-1664 (1989)). The synthesis and physico-chemical properties of partial P-methoxy oligodeoxyribonucleotides were also determined. Only thymidine and cytidine oligomers with 5 methyl phosphotriester could be prepared because of the difficulties encountered in maintaining methyl triester intact. Furthermore, the methyl group was found to have destabilizing effect on the hybridization properties of the modified oligomers with their complementary sequences as compared to the unmodified parent oligodeoxyribonucleotide (Vinogradov, S., Asseline, U., Thoung, N.T., Tet. Let. 34:5899-5902 (1993)).

Other reports have suggested that P-methoxy oligonucleotides are preferable to P-ethoxy as antisense oligonucleotides because of p-methoxy oligonucleotides showed stronger hybridization than methyl phosphonate or P-ethoxy oligonucleotides (van Genderen, M.H.P., et al., Kon. Ned. Akad. van Wetensch. B90:155-159 (1987); van Genderen, M.H.P., et al., Trav. Chim. Pays Bas 108:28-35 (1989)). P-ethoxy oligonucleotides were reported by van Genderen et al. to hybridize poorly to DNA and were thus deemed less suitable for use as antisense oligonucleotides (Moody, H.M., et al., Nucl. Acid Res. 17:4769-4782 (1989)).

P-isopropoxyphosphoramidites have been synthesized from several nucleosides (Stec, W.J., et al., Tet. Let. 26:2191-2194 (1985)), and a few short oligonucleotides containing P-isopropoxyphosphotriesters were synthesized, and hybridization studies were carried out.

United States Patent No. 5,525,719 to Srivastava, S., and Raza, S.K., June 11, 1996, suggests antisense oligonucleotides consisting of 2'-O-Methyl nucleotides linked by phosphodiester and/or P-ethoxy or P-methoxy, phosphotriester moieties.

Presently there are no nucleic acid chemistries nor any chimeras that have been developed that optimally achieve all the features that are needed to provide an effective antisense oligonucleotide i.e. low toxicity, high



WO 98/13526

PCT/US97/17338

specificity, nuclease resistance, ease of synthesis, RNase H compatibility.

### 3. SUMMARY OF THE INVENTION

5 The present invention describes a class of oligonucleotides that has been optimized to target a specific RNA target for RNase H degradation while remaining resistant to nuclease degradation in plasma and within eukaryotic, especially mammalian cells. The oligonucleotides of the  
10 invention contain no naturally occurring 5'→3'-linked nucleotides. Rather, the invention provides oligonucleotides having two types of nucleotides: 2'-deoxyphosphorothioate, which activate RNase H, and 2'-modified nucleotides, which do not. The linkages between the 2'-modified nucleotides can be  
15 phosphodiesters, phosphorothioate or P-ethoxyphosphodiester.

In addition to 5' and 3' ends, the presently described oligonucleotides comprise an RNase H activating region, and a complementarity region that facilitates hybridization to the target sequence. The RNase H-activating region is typically  
20 a contiguous sequence that contains between three and five 2'-deoxyphosphorothioate nucleotides (to activate bacterial RNase H), and typically between about 3 to 12, more typically 5 and 12, and more preferably between about 5 and 10 2'-deoxyphosphorothioate nucleotides to activate eukaryotic,  
25 particularly mammalian, RNase H.

The 5' and 3' ends of the presently described oligonucleotides are protected from exonuclease degradation via the incorporation of modified 5' and 3' terminal bases that are highly nuclease, particularly exonuclease, resistant  
30 and, optionally, by placing a 3' terminal blocking group.

In a preferred embodiment the RNase H activating region, is composed of highly nuclease resistant phosphorothioate nucleotides that is placed at the 5' end of the oligonucleotide.

35 Accordingly, one embodiment of the present invention is a chimeric oligonucleotide comprising an RNase H-activating region of between three and twelve contiguous

WO 98/13526

PCT/US97/17338

2'-deoxyphosphorothioate-linked bases (i.e., phosphorothioate linked 2'-deoxyribonucleotides); a substantially endonuclease resistant complementarity region of between about nine and about fifty 2'-modified bases; a substantially exonuclease resistant 5' terminus; and a substantially exonuclease resistant 3' terminus.

#### 4. DETAILED DESCRIPTION OF THE INVENTION

##### 4.1. THE STRUCTURE OF THE OLIGONUCLEOTIDES

10 An oligonucleotides of the presently invention typically comprise a 5' exonuclease resistant 5' terminal nucleic acid or linkage, a contiguous RNase H activating region of about 3 to about ten bases in length, a 3'-terminal 5'→3'-linked, or optionally 3'-3' linked, e.g., "inverted", nucleoside, and  
15 from about 9 to about 50 5'→3' linked nucleotides, which nucleotides can be 2'-deoxynucleotides or 2'-modified nucleotides that facilitate hybridization of the oligonucleotide to the target mRNA, such as 2'-fluoro, 2'-methoxy, 2'-ethoxy, 2'-methoxyethoxy, 2'-allyloxy (-  
20  $\text{OCH}_2\text{CH}=\text{CH}_2$ ) nucleotides (hereinafter "2'-modified nucleotides"). The 3' terminal nucleoside can, optionally, be a 2'-modified nucleoside. Those skilled in the art appreciate that the 3'-OH of the 3' terminal base can, but need not, be esterified to a phosphate or phosphate analog.  
25 The 3' terminal residue is referred to as a nucleoside even though it may be a nucleotide.

The internucleotide linkages of an oligonucleotide of the invention can be phosphodiester, phosphorothioate or P-ethoxyphosphodiester moieties. The oligonucleotide has a 3'  
30 terminus and a 5' terminus that are substantially protected from nuclease attack. The 3' terminus is protected by having the 3' most 5'→3' linkage or linkages be a phosphorothioate or a P-alkyloxyphosphotriester linkage and/or by having a substituted 3' terminal hydroxyl, e.g., a 3'→3' linked  
35 nucleotide, wherein the alkyloxy radical is methoxy, ethoxy or isopropoxy and, preferably, ethoxy. Preferably two or three 3' terminal internucleotide linkages are

WO 98/13526

PCT/US97/17338

phosphorothioate or a P-alkyloxyphosphotriester linkages. To reduce nuclease degradation, the 5' most 3'→5' linkage preferably should be a phosphorothioate linkage or P-alkyloxyphosphotriester linkage. Preferably, the two 5' most 5 3'→5' linkages should be phosphorothioate linkages or P-ethoxyphosphotriester linkages. Optionally, the 5'-terminal hydroxyl moiety can be esterified with a phosphorus containing moiety, e.g., phosphate, phosphorothioate or P-ethoxyphosphate, without limitation.

- 10 The 3' terminal 5'→3'-linked nucleoside has a 3'-O that can be optionally substituted by a blocking moiety that prevents 3'-exonuclease degradation of the oligonucleotide. In one embodiment, the 3'-hydroxyl is esterified to a nucleotide through a 3'→3' internucleotide linkage.
- 15 Optionally, the 3'→3' linked nucleotide at the 3' terminus can be linked by a phosphorothioate moiety. By incorporating the above chemistries, the presently described oligonucleotides are substantially resistant to 5' and 3' exonucleases and endonucleases. For the purposes of the
- 20 present invention, an oligomer is substantially resistant to a given endo or exonuclease when it is at least about 3-fold more resistant to attack by an endogenous cellular nuclease, and is highly nuclease resistant when it is at least about 6-fold more resistant than a corresponding oligomer comprised
- 25 of unmodified DNA or RNA.

In a preferred embodiment, the oligonucleotide contains, exclusive of an optional blocking nucleotide, between 15 and 50 bases and more preferably between 20 and 30 bases and in a most preferred embodiment the oligonucleotide is 25 bases in

30 length. The oligonucleotide of the invention contains a single contiguous RNase H-activating region of between three to ten 2'-deoxyphosphorothioate nucleotides. The length of the RNase H activating region to activate bacterial RNase H is preferably between three and five nucleotides; to activate

35 a eukaryotic RNase H the activating region is preferably between about five and about ten or twelve nucleotides. The

WO 98/13526

PCT/US97/17338

preferred length of the RNase H-activating region for the  
activation of mammalian RNase H is nine nucleotides.

All 5'→3' linked nucleotides of the oligonucleotide that  
are not a part of the RNase H-activating region are 2'-  
5 modified nucleotides, which contribute to the target binding  
and thus form the complementarity determining region. The  
complementarity region can be a contiguous region or can be  
divided by the RNase H-activating region. In a preferred  
embodiment the complementarity region is a contiguous region,  
10 and, more preferably, is located 3' to the RNase H-activating  
region.

In a preferred embodiment all bases except for the one  
to three 3'-terminal nucleotides and/or nucleoside, the 5'  
terminal nucleotide, and the RNase H activating region  
15 nucleotides, are phosphodiester linked. Large amounts of  
contiguous phosphorothioate linkages are detrimental to the  
function of the oligonucleotides of the invention.  
Accordingly, the oligonucleotides preferably contain not more  
than twelve contiguous phosphorothioate linkages or twelve  
20 contiguous phosphorothioate linked deoxynucleotides.

Additional embodiments of the presently described  
chimeric oligonucleotides have the structures:

5' A:B:C

25 or

5' C:B:A:B:C.

Wherein A is a RNase H activating region of between  
about 3 to about 12 nucleotides, preferably about 3 to about  
10 nucleotides or 5 to about 12 nucleotides long that is also  
30 nuclease stable (e.g., phosphorothioate DNA); B represents a  
region of chemistry (e.g. 2'-O-methyl substituted RNA) that is  
stable against endonucleases (about 4 to about 40 nucleotides  
long; and C represents a one to four nucleotide long  
exonuclease block that typically does not contain  
35 phosphorothioate DNA (i.e., phosphorothioate 2'-O-methyl  
linkages, inverted bases, methylphosphonate, phosphoramidite,  
non-nucleotide linkers, amino linkers, conjugates or any

**WO 98/13526**

PCT/US97/17338

other chemistry consistent with nucleotide synthesis in the art, or yet to be discovered that is not recognized by cellular exonucleases). Alternatively, the configuration may be inverted as follows:

5

5' C:B:A.

If the application does not require activation of RNase H (stearic blocking or triple strand inactivation), the following configuration is useful:

10

5' C:B:C.

#### 4.2. SYNTHESIS OF THE OLIGONUCLEOTIDES

The oligonucleotides of the invention can be synthesized by solid phase or liquid phase nucleotide synthesis, however, synthesis by solid phase techniques is preferred.

Phosphodiester and phosphorothioate linked oligonucleotides can be synthesized, using standard reagents and protocols, on an automated synthesizer utilizing methods that are well known in the art, such as, for example, those disclosed in 20 Stec et al., J. Am. Chem. Soc. 106:6077-6089 (1984); Stec et al., J. Org. Chem. 50(20):3908-3913 (1985); Stec et al., J. Chromatog. 326:263-280 (1985); LaPlanche et al., Nuc. Acid. Res. 14:9081-9093 (1986); and Fasman, G.D., Practical Handbook of Biochemistry and Molecular Biology 1989, CRC

25 Press, Boca Raton, Florida, herein incorporated by reference.

The synthesis of 2'-O-alkyl-oligoribonucleotides, where the alkyl groups are methyl, butyl, allyl or 3,3-dimethylallyl is reviewed by Lamond, Biochem. Soc. Trans. 21:1-8 (1993). Intermediates that are useful in the

30 synthesis of 2'-O-methyl oligoribonucleotides are described  
in U.S. Patents No. 5,013,830, No. 5,525,719 and No.  
5,214,135, which are hereby incorporated by reference.

The synthesis of 2'-fluorophosphodiester and 2'-fluorophosphorothioate oligonucleotides can be performed according to teaching of Kawasaki, A.M., et al., 1993, J. Med. Chem. 36:831-41 and WO 92/03568; the synthesis of P-alkyloxyphosphotriester-linked oligonucleotides and 2'-

WO 98/13526

PCT/US97/17338

modified oligonucleotides can be performed according to U.S. Patent No. 5,525,719, each of which is incorporated herein by reference. The synthesis of phosphorothioate oligodeoxynucleotides is taught by U.S. Patent No. 5,276,019 and No. 5,264,423, which is hereby incorporated by reference. Synthesis of 2'-substituted oligonucleotides can be performed by variations on the techniques disclosed therein.

The synthesis of the oligonucleotides of the invention must be conducted with great attention to quality control.

10 It is particularly important that the phosphorothioate linkages not be contaminated with phosphodiester linkages. It is advisable to pre-test the individual reagent lots to ascertain that high coupling efficiency can be obtained therewith and to exercise all possible precautions to

15 maintain anhydrous conditions.

The quality of the synthesis of oligonucleotides can be verified by testing the oligonucleotides by capillary electrophoresis and denaturing strong anion HPLC (SAX-HPLC). The method of Bergot & Egan, 1992, J. Chrom. 599:35-42 is  
20 suitable. SAX-HPLC is particularly useful to verify that the phosphorothioate nucleotides are completely thiolated, i.e., are not contaminated by a small percentage of phosphodiesters.

The synthesis of oligonucleotides having both  
25 phosphodiester and phosphorothioate linkages is associated with a side reaction whereby the phosphorothioate linkages are oxidized by the standard  $I_2$  treatments that are used to oxidize the cyanoethyl phosphoramidite. This problem can be minimized but not eliminated by reducing the concentration or  
30  $I_2$  to as low as 0.001 M. Therefore, in a preferred embodiment, all phosphorothioates of the oligonucleotides of the invention are found at the 5'-end, so that no phosphorothioate bond is exposed to  $I_2$ .

35 4.3. THE USES OF THE OLIGONUCLEOTIDES

The oligonucleotides of the invention can be used as antisense oligonucleotides in a variety of *in vitro*

WO 98/13526

PCT/US97/17338

experimental situations to specifically degrade an mRNA of unknown function and thereby determine its physiologic function.

The oligonucleotides of the invention can be also used in clinical practice for any disease and against any target RNA for which antisense therapy is now known to be suitable or which is yet to be identified. Medical conditions for which antisense therapy is reported to be suitable includes Respiratory Syncytial Virus infection, WO 95/22553 by Kilkuskie, Influenza Virus infection, WO 94/23028, and malignancies, WO 94/08003. Further examples of clinical uses of antisense oligonucleotides are reviewed, in summary form, in Glaser, V., 1996, Genetic Engineering News 16, 1. Targets of antisense oligonucleotides under that are the subjects of clinical trials include protein kinase C $\alpha$ , ICAM-1, c-raf kinase, p53, c-myb and the bcr/abl fusion gene found in chronic myelogenous leukemia.

## 5. EXAMPLES

### 20 5.1. EXPERIMENTAL CONDITIONS

The antisense activity of the oligonucleotides of the present invention are demonstrated using a test transient expression system which includes an mRNA encoding a luciferase protein that has been modified to include a test sequence derived from the ras gene. The specific antisense effects of an oligonucleotide can be measured by comparing the luciferase production of the test cells with the production of control cells having the same expression plasmid except for the absence of the ras-derived sequence.

25 The oligonucleotides of the invention which were tested have the sequence: 5'-TTGCCCACACCGACGGCGCCACCA-3' (SEQ ID NO: 1)

The details of the assay are as follows:

Plasmid Constructs. The plasmid used for the studies contained a portion of the ras gene sequence fused to luciferase (Monia, B.P., et al. J. Biol. Chem. 267:19954-19962 (1992)). The control luciferase plasmids did not contain the ras target sequence.

WO 98/13526

PCT/US97/17338

Cell Culture Assay. HeLa cells were grown to 40-90% confluence in DMEM/10% FBS, Supplemented with glutamine, penicillin and streptomycin on gelatin coated 24 well plates. The gelatin coating was necessary for cell to remain adherent during the transfections. Prior to transfection the cells were washed twice with PBS (containing magnesium and calcium). LIPOFECTIN™ was mixed gently and 6.6µl was added for each milliliter of reduced serum medium (OPTI-MEM™, Gibco/BRL, Gaithersburg, MD). Oligomers were added from 50-100µM concentrated stock to make a master mixture. The Opti-MEM/LIPOFECTIN/oligomer solution was added to the cells and incubated for 4 hours (~0.5 ml for one well of a 24 well plate).

A target transfection mixture was prepared by first diluting 5µl of lipofectin per ml of OPTI-MEM and mixing. Next 5µg of luciferase target and 5µg of CMV β-galactosidase were added per milliliter of OPTI-MEM/LIPOFECTIN™ mixture. The transfection mixture was mixed gently and allowed to complex for about 15 minutes.. The master mixture reduced error by assuring that the control and experimental cells received the exact same cationic lipid/plasmid complex. The concentration of oligonucleotide in the culture medium was between 200 nM and 400 nM in all experiments. The oligonucleotide containing media was removed from the cells and replaced with growth media and incubated for an additional 9-18 hours. The cells were rinsed with calcium and magnesium free media and the media was removed. The plates were frozen at -70° C for >20 minutes and 100-300 µl of reporter lysis buffer (Promega, Madison WI) was added.

The cells were put through two more freeze thaw cycles, to assure complete lysis. Luciferase assays were performed according to the manufacture's instructions (Promega, Madison WI) and luminescence was detected with a 96 well luminometer (Packard, Meriden CT). β-galactosidase assays were performed (Galacton Plus, Tropix) according to manufactures instructions and detected on the Packard luminometer.



WO 98/13526

PCT/US97/17338

### 5.2. EXPERIMENTAL RESULTS

The results of the luciferase assays are presented in Table I. The results are reported as the percent specific inhibition as calculated by the formula  $100 \times (1 - (LUC_T/LUC_C)^{OLIGO} / (LUC_T/LUC_C)^{NO\ OLIGO})$ ; wherein  $LUC_T$  and  $LUC_C$  are the luciferase levels found in the cells transfected with luciferase plasmids containing and lacking the ras gene insert (SEQ ID NO: 1); and the superscripts "Oligo" and "No Oligo" refer to the presence and absence of antisense oligonucleotides.

TABLE I

	Oligo	Formula	Specific inhibition
15		Controls ("C")	
	C1	25Mo	26%
	C2	25Ms	15%
	C3	9Ds16Mo	15%
	C4	9Do16MoInvT	0%
20	C5	9Dp16MoInvT	18%
	C6	9Dp13Mo3Ms	14%
		Controls with all "S"	
25	S1	25Ds	93%
	S2	16Ms8DsD	100%
	S3	8Ms9Ds7MsM	97%
	S4	9Ds15MsM	95%
30		9Ds at 3' end ("3'I")	
	3'I1	InvTMa15Mo9DsInvT	59%
	3'I2	2Ms14Mo9DsInvT	57%
	3'I3	4Ms12Mo9DsInvT	65%
35		9Ds in Middle ("MI")	
	MI1	5Ms3Mo9Ds4Mo3MsM	64%

WO 98/13526

PCT/US97/17338

	MI2	2Ms6Mo9Ds7 (MsMo) InvT	71%
	MI3	3Ms6Mo9Ds6MoMsInvT	87%
5		9Ds at 5'end ("5'I")	
	5' I1	9Ds16MoInvT	83%
	5' I2	9Ds15MoMsInvT	85%
	5' I3	9Ds16MoBiotin	90%
	5' I4	9Ds16Mp	91%
10	5' I5	9Ds14MoMpD	90%
	5' I6	9Ds13Mo2MpD	94%
	5' I7	9Ds12Mo3MpD	94%
	5' I8	9Ds14MoMsD	93%
	5' I9	9Ds13Mo2MsD	97%
15	5' I10	9Ds12Mo3MsD	95%

Key: M and D refer to 2'-O-methyl- and 2'-deoxy-ribonucleotides, respectively. The letters "o", "s" and "p" refer to phosphodiester, phosphorothioate diester, and P-ethoxy-phosphotriester linked nucleotides. "InvT" refers to a 3'→3' or 5'→5' linked thymidine located at the respective 3' or 5' end of the oligomer.

Table I shows the results of control oligos C1-C6, all phosphorothioate oligos S1-S4, and oligos of the invention having the RNase activating region at the 3' end (3'I1-3'I3), in the middle (MI1-MI3) and at the 5' end (5'I1-5'I10).

Control oligos C1, C2, C5 and C6 showed low levels of specific inhibition because these oligos lacked an RNase H activating region. Oligos C3 and C4 were inactive because the 3' was unprotected and because native ssDNA was unstable, respectively. All phosphorothioate oligonucleotides (S1-S4) showed specific inhibitions that ranged between 93% and 100%, as did oligonucleotides 5'I6-5'I10, which have a 5'-located RNase H activating region and two or three 3' terminal 2'-O-methyl modified P-ethoxy or phosphorothioate linked nucleotides (Mp and Ms, respectively). Lower levels of specific inhibition were observed when oligonucleotides with 3' and mid-located RNase H activating regions were employed or when suboptimal 3' protecting groups were present.

WO 98/13526

PCT/US97/17338

Although the oligonucleotides of the invention having 5' RNase activating regions achieved levels of specific inhibition which were comparable to those achieved by the uniform phosphorothioate oligonucleotides, the oligonucleotides of the invention were superior in that their use was associated with substantially less toxicity. Table II shows specific inhibition, the average metabolic activity as percent of no oligo control, as determined by MTS assay, and the percent viable cells, as determined by trypan blue exclusion for the conventional ("C"), all phosphorothioate ("S"), 3'I, MI and 5'I oligonucleotides; as well as for three species.

TABLE II

Oligo	% INH Luc	% of Control Metabolic Activity	% of Viable Cells
All "O" Oligos C1-C6	15%	94%	76%
All "S" Oligos S1-S4	96%	25%	21%
3'I (1-4)	60%	70%	61%
MI (1-3)	74%	77%	67%
5'I (1-10)	91%	71%	60%

The best oligos on the chart have high percentage values in all columns.

The results demonstrated that the oligonucleotides of the invention achieve levels of specific inhibition more than four times greater than conventional oligonucleotides while showing toxicity levels that were substantially lower than the phosphorothioate oligonucleotides. The optimal group, 5'I, showed specific inhibition that was comparable to the phosphorothioate oligonucleotides.

WO 98/13526

PCT/US97/17338

### 5.3. THE EFFECT OF THE LOCATION OF THE RNase H ACTIVATING REGION

The cause of lower specific activity observed for the 3'I and MI type oligonucleotides was investigated. One possibility was that the oxidation steps using 0.02 M I<sub>2</sub> caused the oxidation of the phosphorothioate linkages to phosphodiester, when phosphodiester linked nucleotides were added 5' to the phosphorothioate linkages. This was found to be the case. Comparison of oligonucleotides 9D<sub>6</sub>15D<sub>0</sub>D ("5'S") and 15D<sub>0</sub>9D<sub>5</sub>D ("3'S") oligonucleotides having the sequence of the test oligonucleotide by analytical HPLC analysis showed that about 85% of the 5'S oligonucleotides were fully thiolated, by contrast only 26% of the 3'S oligonucleotides were completely thiolated (36% were S-1, 24% S-2 and 14% S-3).

Table III shows the distribution of fully thiolated and mono, di and tri-oxidized by-products as a function of the position of the phosphorothiolated region of the oligonucleotide. Four thymidyl pentadodecamers were synthesized using 0.02 M I<sub>2</sub> as the oxidant for 15 nucleotides and a thiolating agent for nine nucleotides.

TABLE III

	Ts	[I <sub>2</sub> ]	S	S-1	S-2	S-3
25	5'-9Ds15DoD03'	0.02M	96%	4%	-	-
	5'-1Do9Ds14DoD-3'	0.02M	85%	15%	-	-
	5'-8Do9Ds7DoD-3'	0.02M	41%	46%	12.5	0.5
	5'-15Do9DsD-3'	0.02M	32%	43%	20%	5%
	5'-15Do9DsD-3'	0.001M	78%	14%	8%	-

The results demonstrated that 96% of the 5'S oligonucleotides are fully thiolated and this percentage steadily decreased as the phosphorothioate region was exposed to more frequent oxidation reactions. When the oxidant concentration was reduced to 0.001M, 78% fully thiolated 3'S 25-T oligonucleotides and about 60% of oligonucleotides having the sequence of the SEQ ID NO: 1 were synthesized.

WO 98/13526

PCT/US97/17338

All publications and patents mentioned in the above specification are herein incorporated by reference. Various modifications and variations of the described method and system of the invention will be apparent to those skilled in the art without departing from the scope and spirit of the invention. Although the invention has been described in connection with specific preferred embodiments, it should be understood that the invention as claimed should not be unduly limited to such specific embodiments. Indeed, various modifications of the above described modes for carrying out the invention which are obvious to those skilled in the field of molecular biology or related fields are intended to be within the scope of the following claims.

15

20

25

30

35

WO 98/13526

PCT/US97/17338

## WE CLAIM:

1. A chimeric antisense oligonucleotide comprising: a 5' terminus; a 3' terminus; and about 11 to about 59 5' to 3' linked nucleotides independently selected from the group consisting of 2'-deoxyphosphorothioate nucleotides, 2'-modified phosphorothioate nucleotides, 2'-modified phosphodiester nucleotides, 2'-modified P-alkoxyphosphotriester nucleotides, wherein:
  - a) said oligonucleotide incorporates an RNase H activating region of between about 3 and about 12 contiguous 2'-deoxyphosphorothioate-linked bases;
  - b) the 5' most 5' to 3' nucleotide linkage is a phosphorothioate or a P-alkoxyphosphodiester linkage;
  - c) the 3' most 5' to 3' nucleotide linkage is a phosphorothioate or P-alkoxyphosphodiester linkage or the 3' terminus is blocked; and
  - d) the oligonucleotide contains not more than 12 contiguous 2'-deoxyphosphorothioate linkages.
2. The oligonucleotide of claim 1, provided the 3' terminus is not blocked by a 3' to 3' phosphorothioate linked nucleotide.
3. The oligonucleotide of claim 1, in which the 3' terminus is blocked by a moiety comprising a 3' to 3' phosphorothioate linked nucleotide.
4. The oligonucleotide of claim 1, in which the 3' terminus is blocked by a moiety comprising a 3' to 3' phosphodiester linked nucleotide.
5. The oligonucleotide of claim 4, in which the 3' most 5' to 3' nucleotide linkage is a phosphorothioate linkage or a P-ethoxyphosphotriester linkage.
6. The oligonucleotide of claim 4, in which the 5' most 5' to 3' nucleotide linkage is a phosphorothioate linkage or a P-ethoxyphosphotriester linkage.

WO 98/13526

PCT/US97/17338

7. The oligonucleotide of claim 1, in which 2'-modified phosphorothioate nucleotides are present at both the 3' terminus and the 5' terminus.
8. The oligonucleotide of claim 1, in which the RNase H activating region is located at the 5' terminus.
9. The oligonucleotide of claim 8, in which the 3' most 5' to 3' nucleotide linkage is a phosphorothioate linkage or a P-ethoxyphosphotriester linkage.
10. The oligonucleotide of claim 9, in which the two 3' most 5'→3' internucleotide linkages are independently either a phosphorothioate linkage or a P-ethoxyphosphotriester linkage.
11. The oligonucleotide of claim 9, in which the RNase H activating region is contiguous with the 3' most 5'→3' nucleotide linkage.
12. The oligonucleotide of claim 11, in which the 2'-modified phosphodiester nucleotide is a 2'-methoxy or 2'-fluoro nucleotide.
13. The oligonucleotide of claim 11, which additionally comprises at least thirteen 2'-methoxy phosphodiester nucleotides.
14. The oligonucleotide of claim 11, having between 15 and 50 nucleotides.
15. The oligonucleotide of claim 14, which additionally comprises at least eight 2'-methoxy phosphodiester nucleotides.
16. The oligonucleotide of claim 14, which additionally comprises at least thirteen 2'-methoxy phosphodiester nucleotides.
17. The oligonucleotide of claim 1, in which the RNase H activating region comprises the 3' terminus.
18. The oligonucleotide of claim 1, in which the 2'-modified phosphodiester nucleotides are selected from the group consisting of 2'-fluoro and 2'-methoxy nucleotides.

WO 98/13526

PCT/US97/17338

19. The oligonucleotide of claim 4, in which the RNase H activating region is present at the 5' terminus followed by four to about forty 5' to 3' linked 2'-methoxy nucleotides, and the 3' terminus is blocked by a 3' to 3' phosphodiester linked deoxyribonucleotide.

20. A method of specifically cleaving an RNA in a cell containing RNase H which comprises administering an effective amount of an oligonucleotide complementary to the RNA, said oligonucleotide comprising: a 5' terminus; a 3' terminus; and about 11 to about 59 5' to 3'-linked nucleotides independently selected from the group consisting of 2'-deoxyphosphorothioate nucleotides, 2'-modified phosphorothioate nucleotides, 2'-modified phosphodiester nucleotides, 2'-modified P-alkyloxyphosphotriester nucleotides, wherein:

a) said oligonucleotide incorporates an RNase H activating region of between about 3 and about 12 contiguous phosphorothioate-linked 2'-deoxynucleotides;

b) the 5' most 5' to 3' nucleotide linkage is a phosphorothioate or a P-alkyloxyphosphodiester linkage;

c) the 3' most 5' to 3' nucleotide linkage is a phosphorothioate or P-alkyloxyphosphodiester linkage or the 3' terminus is blocked; and

d) the oligonucleotide contains not more than twelve contiguous phosphorothioate linked 2'-deoxynucleotides.



## INTERNATIONAL SEARCH REPORT

International application No.  
PCT/US97/17338

<b>A. CLASSIFICATION OF SUBJECT MATTER</b> IPC(6) : C12Q 1/68; C07H 21/04 US CL : 435/6,91.1; 536/24.5 According to International Patent Classification (IPC) or to both national classification and IPC		
<b>B. FIELDS SEARCHED</b> Minimum documentation searched (classification system followed by classification symbols) U.S. : 435/6,91.1; 536/24.5 Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched Electronic data base consulted during the international search (name of data base and, where practicable, search terms used) APS, DIALOG, BIOSIS, MEDLINE, CAS, Derwent Biotechnology Abstracts		
<b>C. DOCUMENTS CONSIDERED TO BE RELEVANT</b>		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	Mesmaeker et al. Backbone modifications in oligonucleotides and peptide nucleic acid systems. Current Opinion in Structural Biology. 1995. Vol. 5. pages 343-355, see entire document.	1-20
<input type="checkbox"/> Further documents are listed in the continuation of Box C. <input type="checkbox"/> See patent family annex.		
* Special categories of cited documents: *A* document defining the general state of the art which is not considered to be of particular relevance *B* earlier document published on or after the international filing date *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) *O* document referring to an oral disclosure, use, exhibition or other means *P* document published prior to the international filing date but later than the priority date claimed *T* later documents published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art *Z* document member of the same patent family		
Date of the actual completion of the international search 17 NOVEMBER 1997		Date of mailing of the international search report 12 DEC 1997
Name and mailing address of the ISA/US Commissioner of Patents and Trademarks Box PCT Washington, D.C. 20231 Facsimile No. (703) 305-3230		Authorized officer John LeGUYADER Telephone No. (703) 308-0196